

## JOINT RANGE OF RÉNYI ENTROPIES

PETER HARREMOËS

The exact range of the joined values of several Rényi entropies is determined. The method is based on topology with special emphasis on the orientation of the objects studied. Like in the case when only two orders of Rényi entropies are studied one can parametrize upper and lower bounds but an explicit formula for a tight upper or lower bound cannot be given.

*Keywords:* Rényi entropies, Shannon entropy, information diagram.

*AMS Subject Classification:* 94A17, 62B10.

## 1. INTRODUCTION

Let  $P = (p_1, p_2, \dots, p_n)$  be a probability vectors. For  $\alpha \in \mathbb{R} \setminus \{0, 1\}$  the Rényi entropy of  $P$  of order  $\alpha$  is defined as a number in  $[0; \infty]$  given by the equation

$$H_\alpha(P) = \frac{1}{1-\alpha} \log \left( \sum_i p_i^\alpha \right).$$

This definition is extended by continuity so that

$$\begin{aligned} H_{-\infty}(P) &= -\log \min_i p_i ; \\ H_0(P) &= \log(\text{number of } p_i \neq 0); \\ H_1(P) &= -\sum_i p_i \log p_i ; \\ H_\infty(P) &= -\log \max_i p_i . \end{aligned}$$

The Rényi entropy  $H_0$  is essentially the Hartley entropy, and was one among other sources of inspiration to Shannon's information theory. The Rényi entropy of order  $\infty$  is also called the min-entropy and essentially related to the "probability of error". The Rényi entropy  $H_2$  is related to index of coincidence and other quantities used for special purposes in crypto analysis, physics etc. [2, 8].

For all  $\alpha$  the Rényi entropy  $H_\alpha$  has the nice property of being additive on product measures. In noiseless source coding for finite systems one wants to avoid very long code words. For such systems the Rényi entropy of some order  $\alpha < 1$  (depending

on the memory of the system) determines how much the source can be compressed. Rényi entropies are also related to general cut-off rates and "guess-work moments" [1, 4].

The relation between  $H_0$  and  $H_1$  is given by the simple inequality

$$H_1(P) \leq H_0(P).$$

This is a special case of the general result that

$$\alpha \rightarrow H_\alpha(P)$$

is a strictly decreasing function except for uniform distributions where it is constant, which follows from a simple application of Jensen's Inequality. The relation between  $H_1$  and  $H_\infty$  has been determined independently in various articles [3, 5, 6, 11, 12]. The relation between Shannon entropy and  $H_2$  has been studied in [7] and in more detail in [8]. The result is illustrated on Figure 1 and by the following theorem.

**Theorem 1** *The upper bound on  $H_2(P)$  given  $H_1(P)$  is attained by a mixture of uniform distributions on  $k$  and  $k+1$  points where  $k$  is determined by the condition  $\log k \leq H_1(P) < \log(k+1)$ . The lower bound on  $H_2(P)$  is attained by a mixture of the uniform distribution on  $n$  points and a uniform distribution on a singleton.*




Figure 1: Range of  $P \rightarrow (H_1(P), H_2(P))$  for a four element set.

In this paper we shall generalize this result and determine the joint range of several Rényi entropies. In general the boundary can be parametrized, but upper and lower bounds cannot be given by explicit formulas. The reason is that the inverse of the function  $s \rightarrow H_\alpha(sU_k + (1-s)U_{k+1})$ , where  $U_k$  and  $U_{k+1}$  are uniform distributions, is in general not an elementary function.

Recently the joint range of Rényi entropies has been used to determine the relative Bahadur efficiency of various power divergence statistics [9,10]. In these papers the joint range of  $H_1$  and  $H_\alpha$  was used with a reference to [8] where the general result for comparison of two Rényi entropies was mentioned without proof. In some cases in physics, joint values of  $H_2(P)$  and  $H_3(P)$  can be measured or computed and one is interested in bounds on  $H_1$  [13]. In order to get bounds on  $H_1$  one is interested in the exact range of the mapping

$$\Psi : P \rightarrow (H_1(P), H_2(P), H_3(P)).$$

In this paper the methods developed in [8] will be refined in order to be able to describe the joint range of in principle any number of Rényi entropies of positive order. We restrict our attention to non-negative orders because these are the most important for applications and because Rényi entropies of negative orders are not continuous near uniform distributions. Although the method is very general we shall only go into details in the cases where two or three Rényi entropies are compared. The main result is that the range has a boundary that can be parametrized by certain mixtures of uniform distributions.

## 2. REDUCTION TO MIXTURES OF UNIFORM DISTRIBUTIONS

A probability vector  $P$  on a set with  $n$  elements can be parametrized by its point probabilities as  $(p_1, p_2, \dots, p_n)$  where  $p_j \geq 0$  and

$$\sum_{j=1}^n p_j = 1.$$

Here we shall assume that  $n$  is fixed so that that  $H_0(P) \leq \log n$ . In order to study the range of  $P \rightsquigarrow (H_{\alpha_1}(P), H_{\alpha_2}(P), \dots, H_{\alpha_m}(P))$  we first consider the related map

$$P \rightarrow \begin{pmatrix} \frac{1}{1-\alpha_1} \log \left( \sum p_j^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( \sum p_j^{\alpha_2} \right) \\ \vdots \\ \frac{1}{1-\alpha_m} \log \left( \sum p_j^{\alpha_m} \right) \\ \sum p_j \end{pmatrix}. \tag{1}$$

The matrix of partial derivatives is

$$\begin{pmatrix} \frac{\alpha_1}{1-\alpha_1} \frac{p_1^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \frac{\alpha_1}{1-\alpha_1} \frac{p_2^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \cdots & \frac{\alpha_1}{1-\alpha_1} \frac{p_{n-1}^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \frac{\alpha_1}{1-\alpha_1} \frac{p_n^{\alpha_1-1}}{\sum p_j^{\alpha_1}} \\ \frac{\alpha_2}{1-\alpha_2} \frac{p_1^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \frac{\alpha_2}{1-\alpha_2} \frac{p_2^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \cdots & \frac{\alpha_2}{1-\alpha_2} \frac{p_{n-1}^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \frac{\alpha_2}{1-\alpha_2} \frac{p_n^{\alpha_2-1}}{\sum p_j^{\alpha_2}} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{\alpha_m}{1-\alpha_m} \frac{p_1^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \frac{\alpha_m}{1-\alpha_m} \frac{p_2^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \cdots & \frac{\alpha_m}{1-\alpha_m} \frac{p_{n-1}^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \frac{\alpha_m}{1-\alpha_m} \frac{p_n^{\alpha_m-1}}{\sum p_j^{\alpha_m}} \\ 1 & 1 & \cdots & 1 & 1 \end{pmatrix}.$$

If this matrix has rank  $m + 1$  in a neighborhood of a point  $P = (p_1, p_2, \dots, p_n)$  then the map (1) is open, i.e. it maps open sets into open sets and a neighborhood of  $P$  is mapped into a neighborhood of the image.

Next we show that if  $P$  has  $m + 1$  different point probabilities then  $P$  is mapped into an interior point in the range. Therefore, assume that  $P$  has  $m + 1$  different point probabilities. For simplicity we may assume that these  $m + 1$  different point probabilities are the first ones and that  $0 < p_1 < p_2 < \dots < p_{m+1}$ . Then

$$\begin{aligned} & \begin{vmatrix} \frac{\alpha_1}{1-\alpha_1} \frac{p_1^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \frac{\alpha_1}{1-\alpha_1} \frac{p_2^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \dots & \frac{\alpha_1}{1-\alpha_1} \frac{p_m^{\alpha_1-1}}{\sum p_j^{\alpha_1}} & \frac{\alpha_1}{1-\alpha_1} \frac{p_{m+1}^{\alpha_1-1}}{\sum p_j^{\alpha_1}} \\ \frac{\alpha_2}{1-\alpha_2} \frac{p_1^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \frac{\alpha_2}{1-\alpha_2} \frac{p_2^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \dots & \frac{\alpha_2}{1-\alpha_2} \frac{p_m^{\alpha_2-1}}{\sum p_j^{\alpha_2}} & \frac{\alpha_2}{1-\alpha_2} \frac{p_{m+1}^{\alpha_2-1}}{\sum p_j^{\alpha_2}} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{\alpha_m}{1-\alpha_m} \frac{p_1^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \frac{\alpha_m}{1-\alpha_m} \frac{p_2^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \dots & \frac{\alpha_m}{1-\alpha_m} \frac{p_m^{\alpha_m-1}}{\sum p_j^{\alpha_m}} & \frac{\alpha_m}{1-\alpha_m} \frac{p_{m+1}^{\alpha_m-1}}{\sum p_j^{\alpha_m}} \\ 1 & 1 & \dots & 1 & 1 \end{vmatrix} \\ & = \left( \prod_{i=1}^m \frac{\alpha_i}{1-\alpha_i} \cdot \prod_{i=1}^m \frac{1}{\sum_j p_j^{\alpha_i}} \right) \begin{vmatrix} p_1^{\alpha_1-1} & p_2^{\alpha_1-1} & \dots & p_m^{\alpha_1-1} & p_{m+1}^{\alpha_1-1} \\ p_1^{\alpha_2-1} & p_2^{\alpha_2-1} & \dots & p_m^{\alpha_2-1} & p_{m+1}^{\alpha_2-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_1^{\alpha_m-1} & p_2^{\alpha_m-1} & \dots & p_m^{\alpha_m-1} & p_{m+1}^{\alpha_m-1} \\ 1 & 1 & \dots & 1 & 1 \end{vmatrix} \end{aligned} \quad (2)$$

Note that the last row can be written as  $(p_1^{\alpha-1} \ p_2^{\alpha-1} \ \dots \ p_m^{\alpha-1} \ p_{m+1}^{\alpha-1})$  with  $\alpha = 1$ . The last determinant is a generalization of the Vandermonde determinant. Like a Vandermonde determinant, it is non-zero if and only if the entries are different, which is the next we have to prove.

**Lemma 2** *Assume that  $0 < x_1 \leq x_2 \leq \dots \leq x_\ell$  and  $\beta_1 < \beta_2 < \dots < \beta_\ell$ . Then the generalized Vandermonde determinant*

$$\det \left( (x_i^{\beta_j})_{i,j=1,2,\dots,\ell} \right) = \begin{vmatrix} x_1^{\beta_1} & x_1^{\beta_2} & \dots & x_1^{\beta_\ell} \\ x_2^{\beta_1} & x_2^{\beta_2} & \dots & x_2^{\beta_\ell} \\ \vdots & \vdots & \ddots & \vdots \\ x_\ell^{\beta_1} & x_\ell^{\beta_2} & \dots & x_\ell^{\beta_\ell} \end{vmatrix}$$

is non-negative. It is zero if and only if there exists  $j \in \{1, 2, \dots, \ell - 1\}$  such that  $x_j = x_{j+1}$ .

**Proof** The proof is by induction in  $\ell$ . For  $\ell = 1$  the generalized Vandermonde determinant is obviously positive. Assume that the result holds for  $\ell = k - 1$ . We have to prove it for  $\ell = k$ . First we have

$$\begin{vmatrix} x_1^{\beta_1} & x_2^{\beta_1} & \dots & x_k^{\beta_1} \\ x_1^{\beta_2} & x_2^{\beta_2} & \dots & x_k^{\beta_2} \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{\beta_k} & x_2^{\beta_k} & \dots & x_k^{\beta_k} \end{vmatrix} = \prod_{j=1}^k x_j^{\beta_1} \begin{vmatrix} 1 & 1 & \dots & 1 \\ x_1^{\beta_2-\beta_1} & x_2^{\beta_2-\beta_1} & \dots & x_k^{\beta_2-\beta_1} \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{\beta_k-\beta_1} & x_2^{\beta_k-\beta_1} & \dots & x_k^{\beta_k-\beta_1} \end{vmatrix}.$$

Therefore without loss of generality we may assume that  $\beta_1 = 0$ . Therefore we have to prove that

$$\begin{vmatrix} 1 & 1 & \cdots & 1 & 1 \\ x_1^{\beta_2} & x_2^{\beta_2} & \cdots & x_{k-1}^{\beta_2} & x_k^{\beta_2} \\ x_1^{\beta_3} & x_2^{\beta_3} & \cdots & x_{k-1}^{\beta_3} & x_k^{\beta_3} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_1^{\beta_k} & x_2^{\beta_k} & \cdots & x_{k-1}^{\beta_k} & x_k^{\beta_k} \end{vmatrix}$$

is non-negative. If  $x_k = x_{k-1}$  the last two columns are identical and determinant is zero so it is sufficient to prove that the partial derivative with respect to  $x_k$  is non-negative. The partial derivative is

$$\begin{vmatrix} 1 & 1 & \cdots & 1 & 0 \\ x_1^{\beta_2} & x_2^{\beta_2} & \cdots & x_3^{\beta_2} & \beta_2 x_k^{\beta_2-1} \\ x_1^{\beta_3} & x_2^{\beta_3} & \cdots & x_3^{\beta_3} & \beta_3 x_k^{\beta_3-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_1^{\beta_k} & x_2^{\beta_k} & \cdots & x_3^{\beta_k} & \beta_k x_k^{\beta_k-1} \end{vmatrix}.$$

Similarly we may take partial derivatives with respect to  $x_{k-1}, x_{k-2}, \dots, x_3$  and  $x_2$  and get

$$\begin{aligned} & \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ x_1^{\beta_2} & \beta_2 x_2^{\beta_2-1} & \beta_2 x_3^{\beta_2-1} & \cdots & \beta_2 x_k^{\beta_2-1} \\ x_1^{\beta_3} & \beta_3 x_2^{\beta_3-1} & \beta_3 x_3^{\beta_3-1} & \cdots & \beta_3 x_k^{\beta_3-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1^{\beta_k} & \beta_k x_2^{\beta_k-1} & \beta_k x_3^{\beta_k-1} & \cdots & \beta_k x_k^{\beta_k-1} \end{vmatrix} \\ &= \begin{vmatrix} \beta_2 x_2^{\beta_2-1} & \beta_2 x_3^{\beta_2-1} & \cdots & \beta_2 x_k^{\beta_2-1} \\ \beta_3 x_2^{\beta_3-1} & \beta_3 x_3^{\beta_3-1} & \cdots & \beta_3 x_k^{\beta_3-1} \\ \vdots & \vdots & \ddots & \vdots \\ \beta_k x_2^{\beta_k-1} & \beta_k x_3^{\beta_k-1} & \cdots & \beta_k x_k^{\beta_k-1} \end{vmatrix} \\ &= \prod_{j=2}^k \beta_j \begin{vmatrix} x_2^{\beta_2-1} & x_3^{\beta_2-1} & \cdots & x_k^{\beta_2-1} \\ x_2^{\beta_3-1} & x_3^{\beta_3-1} & \cdots & x_k^{\beta_3-1} \\ \vdots & \vdots & \ddots & \vdots \\ x_2^{\beta_k-1} & x_3^{\beta_k-1} & \cdots & x_k^{\beta_k-1} \end{vmatrix}. \end{aligned}$$

This is non-negative according to the induction hypothesis.  $\blacksquare$

We see that if  $0 < \alpha_1 < \cdots < \alpha_m < 1$  then the determinant (2) is positive. It is easy to check that this is also the case with the relaxed condition  $0 < \alpha_1 < \cdots < \alpha_m$ .

The Rényi entropies are symmetric in their entries. Therefore we may restrict our attention to probability vectors with increasing entries, i.e.  $0 \leq p_1 \leq p_2 \leq \cdots \leq p_{m+1}$ . The extreme points in the set of ordered probability vectors are the uniform distributions. Let  $U_k$  denote the uniform distribution  $(0, 0, \dots, 0, \frac{1}{k}, \frac{1}{k}, \dots, \frac{1}{k})$ . Let

$k_1, k_2, \dots, k_\ell$  be a sequence of different numbers in  $\{1, 2, \dots, n\}$ . Then the simplex formed by convex combinations of  $U_{k_1}, U_{k_2}, \dots, U_{k_\ell}$  will shall be denoted  $\Delta_{k_1, k_2, \dots, k_\ell}$  and be given an orientation according to the sequence  $U_{k_1}, U_{k_2}, \dots, U_{k_\ell}$ . Observe that if  $k_1 > k_2 > \dots > k_{m+1}$  then the mapping  $\Delta_{k_1, k_2, \dots, k_m} \rightarrow \mathbb{R}^m$  defined by

$$P \rightarrow \begin{pmatrix} \frac{1}{1-\alpha_1} \log \left( \sum p_j^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( \sum p_j^{\alpha_2} \right) \\ \vdots \\ \frac{1}{1-\alpha_m} \log \left( \sum p_j^{\alpha_m} \right) \end{pmatrix}$$

has positive orientation if  $0 < \alpha_1 < \alpha_2 < \dots < \alpha_m$ .

### 3. JOINT RANGE OF TWO RÉNYI ENTROPIES

First we consider distributions on a set with  $n$  elements. We determine the joint range of  $H_{\alpha_1}$  and  $H_{\alpha_2}$  where we assume that  $0 < \alpha_1 < \alpha_2$ . First we shall also assume that  $\alpha_1, \alpha_2 \in ]0; \infty[ \setminus \{1\}$ . Let  $\Phi$  denote the map

$$P \rightarrow \begin{pmatrix} H_{\alpha_1}(P) \\ H_{\alpha_2}(P) \end{pmatrix}.$$

Assume that  $k_1 > k_2 > k_3$ . Then  $\Phi(U_{k_j})$  lies on the diagonal  $\{(x, x) : x \geq 0\}$ , and these points are ordered,

$$H_\alpha(U_{k_1}) > H_\alpha(U_{k_2}) > H_\alpha(U_{k_3})$$

where  $\alpha = \alpha_1$  or  $\alpha = \alpha_2$ . We know that  $H_{\alpha_1}(P) \geq H_{\alpha_2}(P)$  with equality if and only if  $P$  is a uniform distribution. Therefore  $\Phi$  restricted to  $\Delta_{k_1, k_2, k_3}$  must preserve orientation. We know that  $\Phi$  maps inner points of  $\Delta_{k_1, k_2, k_3}$  into inner points of the range of  $\Phi$  so boundary points of the range of  $\Phi$  must have preimages that are boundary points of  $\Delta_{k_1, k_2, k_3}$ . We follow the conventions from homology theory and calculate the boundary with orientation. The boundary of  $\Phi(\Delta_{k_1, k_2, k_3})$  is

$$\begin{aligned} \partial\Phi(\Delta_{k_1, k_2, k_3}) &= \Phi\partial(\Delta_{k_1, k_2, k_3}) \\ &= \Phi(\Delta_{k_2, k_3} - \Delta_{k_1, k_3} + \Delta_{k_1, k_2}) \\ &= \Phi(\Delta_{k_1, k_2} + \Delta_{k_2, k_3} + \Delta_{k_3, k_1}), \end{aligned}$$

which is just another way of writing the closed curve from  $U_{k_1}$  to  $U_{k_2}$  to  $U_{k_3}$  and back to  $U_{k_1}$ . Therefore any point on the boundary of the range of  $\Phi$  must be the image of a mixture of two uniform distributions.

Assume that  $k_1 > k_2 > k_3 > k_4$ . Then the simplices  $\Delta_{k_1, k_2, k_3}$  and  $\Delta_{k_1, k_3, k_4}$  are

both positively oriented and

$$\begin{aligned}
 \partial\Phi(\Delta_{k_1,k_2,k_3} + \Delta_{k_1,k_3,k_4}) &= \Phi\partial(\Delta_{k_1,k_2,k_3} + \Delta_{k_1,k_3,k_4}) \\
 &= \Phi\left(\begin{array}{c} \partial\Delta_{k_1,k_2,k_3} \\ +\partial\Delta_{k_1,k_3,k_4} \end{array}\right) \\
 &= \Phi\left(\begin{array}{c} \Delta_{k_2,k_3} - \Delta_{k_1,k_3} + \Delta_{k_1,k_2} \\ +\Delta_{k_3,k_4} - \Delta_{k_1,k_4} + \Delta_{k_3,k_4} \end{array}\right) \\
 &= \Phi\left(\begin{array}{c} \Delta_{k_2,k_3} - \Delta_{k_1,k_3} + \Delta_{k_1,k_2} \\ +\Delta_{k_3,k_4} - \Delta_{k_1,k_4} + \Delta_{k_1,k_3} \end{array}\right) \\
 &= \Phi(\Delta_{k_1,k_2} + \Delta_{k_2,k_3} + \Delta_{k_3,k_4} + \Delta_{k_4,k_1}).
 \end{aligned}$$

We see that  $\Phi(\Delta_{k_1,k_3})$  does not contribute to the boundary of

$$\partial\Phi(\Delta_{k_1,k_2,k_3} + \Delta_{k_1,k_3,k_4}).$$

Similarly  $\Phi(\Delta_{k_2,k_4})$  does not contribute to the boundary. We may formulate this result as  $\Delta_{a,b}$  does not contribute to the range if it is a diagonal in a quadruple. The non-diagonal simplices are  $\Delta_{n,n-1}, \Delta_{n-1,n-2}, \dots, \Delta_{2,1}$  and  $\Delta_{1,n}$ . These form a closed curve

$$\Delta_{n,n-1} + \Delta_{n-1,n-2} + \dots + \Delta_{2,1} + \Delta_{1,n}$$

and the boundary is the image of this curve, i.e.

$$\Phi(\Delta_{n,n-1} + \Delta_{n-1,n-2} + \dots + \Delta_{2,1} + \Delta_{1,n}),$$

This result easily extends to the cases where one or more of the orders equal 1 or  $\infty$ . The upper bound does not depend on  $n$  so we get the following theorem.

**Theorem 3** *Assume  $0 < \alpha_1 < \alpha_2$ . Then the upper bound on  $H_{\alpha_2}(P)$  given  $H_{\alpha_1}(P)$  is attained by a mixture of uniform distributions on  $k$  and  $k+1$  points where  $k$  is determined by the condition  $\log k \leq H_{\alpha_1}(P) < \log(k+1)$ .*

For distributions on set with  $n$  elements we also get a tight lower bound, but if we have no restriction on  $n$  the situation is a little more complicated.

**Theorem 4** *Assume  $0 < \alpha_1 < \alpha_2$ . If  $P$  is a distribution on a set with  $n$  elements and  $H_{\alpha_1}(P)$  is fixed then a lower bound on  $H_{\alpha_2}$  is attained for a mixture of the uniform distributions  $U_1$  and  $U_n$ . If no restriction on  $n$  is given and if  $H_{\alpha_1}(P) > 0$  is fixed then a tight lower bound on  $H_{\alpha_2}(P)$  is given by*

$$H_{\alpha_2}(P) > \begin{cases} 0, & \text{if } \alpha_1 \leq 1; \\ \frac{\alpha_2 - \alpha_1 - 1}{\alpha_2 - 1} H_{\alpha_1}(P), & \text{if } \alpha_1 > 1. \end{cases}$$

**Proof** If we have no restriction on  $n$  then the range is

$$\bigoplus_{n=2}^{\infty} \Phi(\Delta_{n+1,n,1}).$$

So we just have to determine the asymptotics of  $\Phi(\Delta_{n,1})$ . The curve  $\Delta_{1,n}$  has the parametrization  $P_t = (\frac{t}{n}, \frac{t}{n}, \dots, \frac{t}{n}, \frac{t}{n} + 1 - t)$ ,  $t \in [0; 1]$ . Therefore the curve  $\Phi(\Delta_{n,1})$  has the parametrization

$$\left( \begin{array}{c} \frac{1}{1-\alpha_1} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_1} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_2} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_2} \right) \end{array} \right).$$

We have to study the asymptotics of this curve for  $n$  tending to infinity. There are several cases and they need separate analysis.

**Case  $\alpha_1 > 1$ .** We also have  $\alpha_2 > 1$  so for a fixed value of  $t$  we get

$$\left( \begin{array}{c} \frac{1}{1-\alpha_1} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_1} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_2} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_2} \right) \end{array} \right) \rightarrow \left( \begin{array}{c} \frac{\alpha_1}{1-\alpha_1} \log(1-t) \\ \frac{\alpha_2}{1-\alpha_2} \log(1-t) \end{array} \right)$$

for  $n$  tending to infinity. Hence the straight line with slope  $\frac{\alpha_2 - \alpha_1 - 1}{\alpha_2 - 1} \frac{\alpha_1 - 1}{\alpha_1}$  is the boundary of the range.

**Case  $\alpha_2 \geq 1$  and  $\alpha_1 \leq 1$ .** First we assume that  $\alpha_1 < 1$ . For a fixed value of the parameter  $t$  the Rényi entropy  $H_{\alpha_2}$  tends to a constant as above but  $H_{\alpha_1}$  tends to infinity. For a fixed value of  $H_{\alpha_1}(P) > 0$  the lower bound  $H_{\alpha_2}(P) > 0$  is tight. This bound is also tight for  $\alpha_1 = 1$  and can be obtained by letting  $\alpha_1$  tend to 1 from above or below.

**Case  $0 < \alpha_2 \leq 1$ .** First assume that  $\alpha_2 < 1$ . If  $t = n^{1-1/\alpha_2}$  then

$$\begin{aligned} & \left( \begin{array}{c} \frac{1}{1-\alpha_1} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_1} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( (n-1) \left( \frac{t}{n} \right)^{\alpha_2} + \left( \frac{t}{n} + 1 - t \right)^{\alpha_2} \right) \end{array} \right) \\ &= \left( \begin{array}{c} \frac{1}{1-\alpha_1} \log \left( n^{-\frac{\alpha_1}{\alpha_2}} \cdot \frac{n-1}{n} + \left( n^{-1/\alpha_2} + 1 - n^{1-1/\alpha_2} \right)^{\alpha_1} \right) \\ \frac{1}{1-\alpha_2} \log \left( \frac{n-1}{n} + \left( n^{-1/\alpha_2} + 1 - n^{1-1/\alpha_2} \right)^{\alpha_2} \right) \end{array} \right). \end{aligned}$$

We see that the second coordinate tends to  $\frac{1}{1-\alpha_2} \log 2$ , while the first coordinate tends to  $\infty$ . Therefore for a fixed value of  $H_{\alpha_1}(P) > 0$  the lower bound  $H_{\alpha_2}(P) > 0$  is tight. Tightness of this bound also holds for  $\alpha_2 = 1$ , which can be seen by letting  $\alpha_2$  tend 1 from above or from below. ■

#### 4. JOINT RANGE OF THREE RÉNYI ENTROPIES

Determining the range of three Rényi entropies is done in the same way as in the previous section. We consider the map  $\Psi$  given by

$$P \rightarrow \left( \begin{array}{c} H_{\alpha_1}(P) \\ H_{\alpha_2}(P) \\ H_{\alpha_3}(P) \end{array} \right).$$

First we consider the situation where the domain consist of distributions on  $n$  points. Boundary points of  $\Psi$  must be images of mixtures of three uniform distributions. If



$n > m > \ell > k > 1$  then the restriction of  $\Psi$  to the simplices  $\Delta_{n,m,\ell,k}$  or to  $\Delta_{m,\ell,k,1}$  conserves orientation. Therefore

$$\begin{aligned} \partial\Psi(\Delta_{n,m,\ell,k} + \Delta_{m,\ell,k,1}) &= \partial\Psi(\partial\Delta_{n,m,\ell,k} + \partial\Delta_{m,\ell,k,1}) \\ &= \partial\Psi\left(\begin{array}{c} \Delta_{m,\ell,k} - \Delta_{n,\ell,k} + \Delta_{n,m,k} - \Delta_{n,m,\ell} \\ +\Delta_{\ell,k,1} - \Delta_{m,k,1} + \Delta_{m,\ell,1} - \Delta_{m,\ell,k} \end{array}\right) \\ &= \partial\Psi\left(\begin{array}{c} -\Delta_{n,\ell,k} + \Delta_{n,m,k} - \Delta_{n,m,\ell} \\ +\Delta_{\ell,k,1} - \Delta_{m,k,1} + \Delta_{m,\ell,1} \end{array}\right). \end{aligned}$$

We see that  $\Delta_{m,\ell,k}$  gives no contribution to the boundary and therefore only simplices  $\Delta_{m,\ell,k}$  with either  $m = n$  or  $k = 1$  give a contributions to the boundary.

If  $n > m > \ell > k > 1$  then the restriction of  $\Psi$  to the simplices  $\Delta_{n,m,k,1}$  or to  $\Delta_{m,\ell,k,1}$  conserves orientation. Therefore

$$\begin{aligned} \partial\Psi(\Delta_{n,m,k,1} + \Delta_{m,\ell,k,1}) &= \partial\Psi(\partial\Delta_{n,m,k,1} + \partial\Delta_{m,\ell,k,1}) \\ &= \partial\Psi\left(\begin{array}{c} \Delta_{m,k,1} - \Delta_{n,k,1} + \Delta_{n,m,1} - \Delta_{n,m,k} \\ +\Delta_{\ell,k,1} - \Delta_{m,k,1} + \Delta_{m,\ell,1} - \Delta_{m,\ell,k} \end{array}\right) \\ &= \partial\Psi\left(\begin{array}{c} -\Delta_{n,k,1} + \Delta_{n,m,1} - \Delta_{n,m,k} \\ +\Delta_{\ell,k,1} + \Delta_{m,\ell,1} - \Delta_{m,\ell,k} \end{array}\right). \end{aligned}$$

We see that the simplex  $\Delta_{m,k,1}$  gives no contribution to the boundary of the range of  $\Psi$ . So if  $m < n$  the simplex  $\Delta_{m,k,1}$  can only give a contribution to the boundary if there exist no natural number  $\ell$  such that  $m > \ell > k$ , i.e.  $k = m - 1$ . In the same way we can show that a simplex of the form  $\Delta_{n,m,\ell}$  will only contribute to the boundary if  $\ell = m - 1$  and that a simplex  $\Delta_{n,m,1}$  only contributes if  $m = n - 1$  or if  $m = 2$ . Thus the boundary of the range consist of images of the simplices  $\Delta_{m,m-1,1}$  and of the form  $\Delta_{n,m,m-1}$ . Here we notice that

$$\begin{aligned} \partial\left(\bigoplus_{m=3}^n \Delta_{m,m-1,1} - \bigoplus_{m=2}^{n-1} \Delta_{n,m,m-1}\right) &= \bigoplus_{m=3}^n \partial\Delta_{m,m-1,1} - \bigoplus_{m=2}^{n-1} \partial\Delta_{n,m,m-1} \\ &= \bigoplus_{m=3}^n (\Delta_{m-1,1} - \Delta_{m,1} + \Delta_{m,m-1}) \\ &\quad - \bigoplus_{m=2}^{n-1} (\Delta_{m,m-1} - \Delta_{n,m-1} + \Delta_{n,m}) = 0, \end{aligned}$$

so that

$$\bigoplus_{m=3}^n \Delta_{m,m-1,1} - \bigoplus_{m=2}^{n-1} \Delta_{n,m,m-1}$$

is a closed surface and that the range of  $\Psi$  has the image of this surface as boundary.

It is possible to describe the situation in more detail. Let  $\Phi$  denote the map

$$P \rightarrow \begin{pmatrix} H_{\alpha_1}(P) \\ H_{\alpha_2}(P) \end{pmatrix}.$$




Figure 2: The left diagram illustrates the range of  $\Delta_{4,2,1}$  and  $\Delta_{4,3,2}$  with orientation ( $n = 4$ ). The range  $\Psi$  applied to these simplices give lower bounds on  $H_3$ . The right diagram illustrates the range of  $\Delta_{3,1,2}$  and  $\Delta_{4,1,3}$  with orientation ( $n = 4$ ). The range  $\Psi$  applied to these simplices give upper bounds on  $H_3$ .

Then  $\Phi$  restricted to  $\bigoplus_{m=3}^n \Delta_{m,m-1,1}$  is a homeomorphism. If

$$\Phi(P) = \begin{pmatrix} a \\ b \end{pmatrix}$$

then there exist a unique  $m$  and unique weights  $x, y, z \geq 0$  that sum up to 1 such that  $P = x \cdot U_m + y \cdot U_{m-1} + z \cdot U_1$ . For any distribution  $Q$  with  $\Phi(Q) = \begin{pmatrix} a \\ b \end{pmatrix}$  we have  $H_{\alpha_3}(Q) \leq H_{\alpha_3}(P)$ . Thus,  $\bigoplus_{m=3}^n \Delta_{m,m-1,1}$  gives tight upper bounds on  $H_{\alpha_3}$  in terms of  $H_{\alpha_1}$  and  $H_{\alpha_2}$ . We notice that this upper bound does not depend on  $n$ . Similarly, the lower bound on  $H_{\alpha_3}$  for fixed  $H_{\alpha_1}$  and  $H_{\alpha_2}$  is determined by the surface  $\bigoplus_{m=2}^{n-1} \Delta_{n,m,m-1}$  and just as in the case of two Rényi entropies the lower bound will depend on  $n$ .

## 5. DISCUSSION

The result can be seen as a generalization of the result in [8]. The essential step in the whole construction is the positivity of the generalized Vandermonde determinant. Therefore the construction can be iterated so that one in principle can determine the boundary of the range of any number of Rényi entropies of positive order.

## ACKNOWLEDGEMENT

I thank Karol Zyczkowski for useful discussions. His paper [13] was an important inspiration for this article. He, Flemming Topsøe and Christian Schaffner have also contributed with several important remarks to this paper.

The author was supported by grants from Villum Kann Rasmussen Foundation, The Banach Center, INTAS (project 00-738), Danish Natural Research Council, and the European Pascal Network of Excellence.

(Received April xx, 2009.)

---

#### REFERENCES

---

- [1] E. Arikan. An inequality on guessing and its application to sequential decoding. *IEEE Trans. Inform. Theory*, 42(1):99–105, Jan. 1996.
- [2] C. Arndt. *Information Measures*. Springer, Berlin, 2001.
- [3] M. Ben-Bassat.  $f$ -entropies, probability of error, and feature selection. *Information and Control*, 39:227–242, 1978.
- [4] I. Csiszár. Generalized cutoff rates and Rényi information measures. *IEEE Trans. Inform. Theory*, 41(1):26–34, Jan. 1995.
- [5] M. Feder and N. Merhav. Relations between entropy and error probability. *IEEE Trans. Inform. Theory*, 40:259–266, 1994.
- [6] J. D. Golić. On the relationship between the information measures and the Bayes probability of error. *IEEE Trans. Inform. Theory*, 33(5):681–690, Sept. 1987.
- [7] A. György and T. Linder. Optimal entropy-constrained scalar quantization of a uniform source. *IEEE Trans. Inform. Theory*, IT-46(7):2704–2711, Nov. 2000.
- [8] P. Harremoës and F. Topsøe. Inequalities between entropy and index of coincidence derived from information diagrams. *IEEE Trans. Inform. Theory*, 47(7):2944–2960, Nov. 2001.
- [9] P. Harremoës and I. Vajda. Efficiency of entropy testing. In *International Symposium on Information Theory*, pages 2639–2643. IEEE, July 2008.
- [10] P. Harremoës and I. Vajda. On the Bahadur-efficient testing of uniformity by means of the entropy. *IEEE Trans. Inform. Theory*, 54(1):321–331, Jan. 2008.
- [11] V. A. Kovalevskij. *The problem of character recognition from the point of view of mathematical statistics*, pages 3–30. Spartan, New York, 1967.
- [12] D. L. Tebbe and S. J. Dwyer. Uncertainty and the probability of error. *IEEE Trans. Inform. Theory*, 14:516–518, 1968.
- [13] K. Zyczkowski. Rényi extrapolation of Shannon entropy. *Open Systems and Information Dynamics*, 10:297–310, 2003.

P. Harremoës, Centrum Wiskunde & Informatica, Science Park 123, 1090 GB Amsterdam, Noord-Holland, The Netherlands.

E-mail: P.Harremoës@cwi.nl